



The American Heart Association and the American Medical Association recognize these practices



TARGET: BP



IOWA

KANSAS

KENTUCKY

LOUISIANA

MARYLAND

the 1990s, the incidence of *S. flexneri* infections in the United Kingdom has increased, and the incidence of *S. flexneri* infection in the United States has increased in the 1980s and 1990s [10, 11].

There is a paucity of data on the incidence of *S. flexneri* infection in the United Kingdom. The only published data on the incidence of *S. flexneri* infection in the United Kingdom are from a study of 10 years of *S. flexneri* infections in the United Kingdom, which reported an incidence of 1.5 cases per 100 000 per year [12].

The purpose of this study was to determine the incidence of *S. flexneri* infection in the United Kingdom, and to determine the risk factors for *S. flexneri* infection. The study was conducted in the United Kingdom, and the results are presented in this paper.

METHODS

Study area

The study was conducted in the United Kingdom, and the results are presented in this paper. The study was conducted in the United Kingdom, and the results are presented in this paper.

The study was conducted in the United Kingdom, and the results are presented in this paper. The study was conducted in the United Kingdom, and the results are presented in this paper.

The study was conducted in the United Kingdom, and the results are presented in this paper. The study was conducted in the United Kingdom, and the results are presented in this paper.

The study was conducted in the United Kingdom, and the results are presented in this paper. The study was conducted in the United Kingdom, and the results are presented in this paper.

The study was conducted in the United Kingdom, and the results are presented in this paper. The study was conducted in the United Kingdom, and the results are presented in this paper.

The study was conducted in the United Kingdom, and the results are presented in this paper. The study was conducted in the United Kingdom, and the results are presented in this paper.

The study was conducted in the United Kingdom, and the results are presented in this paper. The study was conducted in the United Kingdom, and the results are presented in this paper.

the 1990s, the incidence of *S. flexneri* infections in the United Kingdom has increased, and the incidence of *S. flexneri* infection in the United States has increased in the 1980s and 1990s [10, 11].

There is a paucity of data on the incidence of *S. flexneri* infection in the United Kingdom. The only published data on the incidence of *S. flexneri* infection in the United Kingdom are from a study of 10 years of *S. flexneri* infections in the United Kingdom, which reported an incidence of 1.5 cases per 100 000 per year [12].

The purpose of this study was to determine the incidence of *S. flexneri* infection in the United Kingdom, and to determine the risk factors for *S. flexneri* infection. The study was conducted in the United Kingdom, and the results are presented in this paper.

METHODS

Study area

The study was conducted in the United Kingdom, and the results are presented in this paper. The study was conducted in the United Kingdom, and the results are presented in this paper.

The study was conducted in the United Kingdom, and the results are presented in this paper. The study was conducted in the United Kingdom, and the results are presented in this paper.

The study was conducted in the United Kingdom, and the results are presented in this paper. The study was conducted in the United Kingdom, and the results are presented in this paper.

The study was conducted in the United Kingdom, and the results are presented in this paper. The study was conducted in the United Kingdom, and the results are presented in this paper.

The study was conducted in the United Kingdom, and the results are presented in this paper. The study was conducted in the United Kingdom, and the results are presented in this paper.

The study was conducted in the United Kingdom, and the results are presented in this paper. The study was conducted in the United Kingdom, and the results are presented in this paper.

The study was conducted in the United Kingdom, and the results are presented in this paper. The study was conducted in the United Kingdom, and the results are presented in this paper.

the 1990s, the incidence of *S. flexneri* infections in the United Kingdom has increased, and the incidence of *S. flexneri* infection in the United States has increased in the 1980s and 1990s [10, 11].

There is a paucity of data on the incidence of *S. flexneri* infection in the United Kingdom. The only published data on the incidence of *S. flexneri* infection in the United Kingdom are from a study of 10 years of *S. flexneri* infections in the United Kingdom, which reported an incidence of 1.5 cases per 100 000 per year [12].

The purpose of this study was to determine the incidence of *S. flexneri* infection in the United Kingdom, and to determine the risk factors for *S. flexneri* infection. The study was conducted in the United Kingdom, and the results are presented in this paper.

METHODS

Study area

The study was conducted in the United Kingdom, and the results are presented in this paper. The study was conducted in the United Kingdom, and the results are presented in this paper.

The study was conducted in the United Kingdom, and the results are presented in this paper. The study was conducted in the United Kingdom, and the results are presented in this paper.


The study was conducted in the United Kingdom, and the results are presented in this paper. The study was conducted in the United Kingdom, and the results are presented in this paper.


The study was conducted in the United Kingdom, and the results are presented in this paper. The study was conducted in the United Kingdom, and the results are presented in this paper.

The study was conducted in the United Kingdom, and the results are presented in this paper. The study was conducted in the United Kingdom, and the results are presented in this paper.

The study was conducted in the United Kingdom, and the results are presented in this paper. The study was conducted in the United Kingdom, and the results are presented in this paper.

The study was conducted in the United Kingdom, and the results are presented in this paper. The study was conducted in the United Kingdom, and the results are presented in this paper.





MASSACHUSETTS

INDIANA

MICHIGAN

IOWA

KENTUCKY

LOUISIANA

MARYLAND

NORTH DAKOTA

OHIO

Montefiore Bronx Health Collective

OKLAHOMA

NORTH CAROLINA

PUERTO RICO

SOUTH CAROLINA

TENNESSEE

TEXAS

VIRGINIA

Vancouver Clinic - Ridgefield Clinic

WASHINGTON

