

RECOGNITION MEASURES

- **Diabetes Treatment** - Percent of diabetic patients or newly-diagnosed diabetics receiving diabetes treatment in the form of glycemic control (diet or antihyperglycemic medication) or follow up appointment for diabetes management scheduled at discharge.
- **Lipid Lowering Medications at Discharge for Patients with Diabetes** - Percent of heart failure patients with diabetes who were prescribed lipid lowering medications at discharge.
- **Smoking Cessation Advice for Patients with Diabetes** - Percent of heart failure patients with diabetes and a history of smoking cigarettes, who are given smoking cessation advice or counseling during hospital stay.
- **SGLT-2 Inhibitor for Patients with Diabetes** - Percent of patients with heart failure and type 2 (sodium glucose cotransporter-2 or SGLT-2 Inhibitor).

Hospitals must achieve 80% compliance for 12 consecutive months for a composite of the below measures::

- **IV Alteplase Arrive by 3.5 Hour, Treat by 4.5 Hour (Patients with Diabetes)** - Percent of acute ischemic stroke patients with diabetes who arrive at the hospital within 210 minutes (3.5 hours) of time last known well and for whom IV thrombolytic was initiated at this hospital within 270 minutes (4.5 hours) of time last known well.
- **Early Antithrombotics for Patients with Diabetes** - Percent of patients with ischemic stroke or TIA and diabetes who receive antithrombotic therapy by the end of hospital day two.
- **VTE Prophylaxis for Patients with Diabetes** - Percent of patients with diabetes and an ischemic stroke, or a hospital admission.
- **Antithrombotics for Patients with Diabetes** - Percent of patients with an ischemic stroke or TIA and diabetes prescribed antithrombotic therapy at discharge.
- **Anticoagulant for AFib/AFlutter for Patients with Diabetes** - Percent of patients with an ischemic stroke
- **Smoking Cessation Advice for Patients with Diabetes** - Percent of patients with ischemic or hemorrhagic stroke or TIA and diabetes and a history of smoking cigarettes, who are, or whose caregivers are, given smoking cessation advice or counseling during hospital stay.
- **Intensive Statin Prescribed at Discharge for Patients with Diabetes** - Percent of ischemic stroke or TIA patients with diabetes who are discharged on statin medication.
- **Diabetes Treatment** - Percent of diabetic patients or newly-diagnosed diabetics receiving diabetes treatment in the form of glycemic control (diet or medication or follow up appointment for diabetes management scheduled at discharge).
- **Therapeutic Lifestyle Recommendation for Patients with Diabetes** - Percent of ischemic stroke or TIA patients with diabetes who receive therapeutic lifestyle recommendations at discharge.
- **SGLT-2 Inhibitor for Patients with Diabetes** - Percent of ischemic stroke or TIA patients with type 2 (glucagon-like peptide-1 receptor/GLP-1 Receptor Agonist or sodium glucose cotransporter 2/SGLT-2 Inhibitor).

